arXiv:hep-ph/0005254v1 25 May 2000

Prompt J/ψ Polarization at the Tevatron

Jungil Lee[a]
II. Institute of Theoretical Physics,
University of Hamburg, 22761 Hamburg, Germany
E-mail: jungil@mail.desy.de

The polarization of prompt J/ψ at the Tevatron is calculated within the nonrelativistic QCD (NRQCD) factorization framework. The contribution from radiative decays of P-wave charmonium states decreases, but does not eliminate, the transverse polarization at large transverse momentum. The prediction agrees with measurements from the CDF collaboration at intermediate values of p_T, but disagrees at the large values of p_T measured.

The NRQCD factorization approach to inclusive quarkonium production[1] makes the remarkable prediction that in hadron collisions $J^{PC} = 1^{--}$ quarkonium states should be transversely polarized at large p_T[2]. In high-energy $p\bar{p}$ collisions, the dominant contribution to the charmonium production rate at large p_T comes from gluon *fragmentation*[3]. At leading order in α_s, a $Q\overline{Q}$ pair with small relative momentum created by the virtual gluon is in a color-octet ${}^3S_1$ state with the same transverse polarization as the almost on-shell gluon. Due to the approximate heavy quark spin symmetry of NRQCD, 1^{--} quarkonium states from the pair should have a large transverse polarization at sufficiently large p_T. Recent measurements at the Tevatron by the CDF collaboration seem to be in contradiction with this prediction[4].

A convenient measure of the polarization is $\alpha = (T-2L)/(T+2L)$, where T and L are the transverse and longitudinal polarization fractions in the hadron CM frame. The variable α describes the angular distribution of leptons from the decay of the J/ψ with respect to the J/ψ momentum. Beneke and Rothstein studied the dominant fragmentation mechanisms for producing longitudinally polarized 1^{--} states[5]. For charmonium production at the Tevatron, one should also take into account the *fusion* contributions from parton processes $ij \to c\bar{c} + k$ since fragmentation does not yet dominate for most of the p_T region. The polarization variable α for direct J/ψ and direct ψ' at the Tevatron have been predicted by Beneke and Krämer[6] and by Leibovich[7]. They predicted that α should be small for p_T below about 5 GeV, but then should begin to rise dramatically. On the contrary, the CDF data shows no sign of transverse polarization of direct ψ' at large p_T[4].

The CDF collaboration has also measured the polarization of *prompt* J/ψ[4]

---

[a] Talk given at 5th Workshop on QCD, 3-7 January 2000, Villefranche-sur-Mer, France.

(*i.e.* J/ψ's that do not come from the decay of B hadrons). The advantage of this measurement is that the number of J/ψ events is larger than for ψ' by a factor of about 100. On the other hand, theoretical predictions of the polarization of prompt J/ψ are complicated since the prompt signal includes J/ψ's that come from decays of the higher charmonium states χ_{c1}, χ_{c2}, and ψ' [8,9]. The polarization of J/ψ from ψ' not via χ_c is straightforward to calculate, since the spin is unchanged by the transition. The polarization of J/ψ from χ_c and of J/ψ from ψ' via χ_c is more complicated, because the χ_{cJ}'s are produced in various spin states and they decay into J/ψ through radiative transitions [10]. Therefore, it is interesting to investigate the cascade effect in this problem.

The *NRQCD factorization formula* for the differential cross section for the inclusive production of a charmonium state H of momentum P and spin quantum number λ has the schematic form

$$d\sigma^{H_\lambda(P)} \;=\; d\sigma^{c\bar{c}_n(P)}\, \langle O_n^{H_\lambda(P)} \rangle, \tag{1}$$

where the implied sum on n extends over all the color and angular momentum states of the $c\bar{c}$ pair. The $c\bar{c}$ cross sections $d\sigma^{c\bar{c}_n}$, which are independent of H, can be calculated using perturbative QCD. All dependence on the state H is contained within the nonperturbative NRQCD matrix elements(ME's) $\langle O_n^{H_\lambda(P)} \rangle$. The Lorentz indices, which are suppressed in (1), are contracted with those of $d\sigma^{c\bar{c}_n}$ to give a scalar cross section. The symmetries of NRQCD can be used to reduce the tensor ME's $\langle O_n^{H_\lambda(P)} \rangle$ to scalar ME's $\langle O_n^H \rangle$ that are independent of P and λ. One may calculate the cross section for polarized quarkonium once the relevant scalar ME's are known.

In $p\bar{p}$ collisions, the parton processes that dominate the $c\bar{c}$ cross section depend on p_T. If p_T is of order m_c, the *fusion* processes dominate. These include the parton processes $ij \to c\bar{c} + k$, with $i, j = g, q, \bar{q}$ and $q = u, d, s$. At p_T much larger than m_c, the parton cross sections are dominated by *fragmentation* processes. To get a consistent LO evaluation of polarization parameter α in this region, one should take care. Since we are interested in the LO prediction of α, we should first get a LO prediction of longitudinal fraction $L/(T+L)$. The LO contributions to L are order-α_s^2 processes $g \to c\bar{c} + g$, while the only contribution to T is the order-α_s process $g \to c\bar{c}_8(^3S_1)$ [5]. Inclusion of the longitudinal part in the fragmentation process is crucial in the calculation of α even though it has only a small contribution to the total cross section. The fragmentation function is evolved using the standard homogeneous timelike evolution equation. Since $c\bar{c}_8(^3S_1) \to \psi_\lambda(nS)$ is included in both the fragmentation and the fusion process, we interpolate between the fusion cross section at low p_T and the fragmentation cross section at high p_T [10,11]. In all the other $c\bar{c} \to H$ channels, we use the fusion cross section.

The color-singlet ME's can be determined phenomenologically from the decay rates for $\psi(nS) \to \ell^+\ell^-$ and $\chi_{c2} \to \gamma\gamma$ [12]. The color-octet ME's are phenomenological parameters and they are determined from production data at the Tevatron [13,8]. In case of χ_c, we have a good agreement with the CDF data on the ratio χ_{c1}/χ_{c2} [9]. More details on the analysis methods are explained in Refs. [10,14]. There are numerous theoretical uncertainties in the polarization calculation. In our analysis, we allow the variations in PDF (CTEQ5L and MRST98LO) [15], factorization and fragmentation scales ($\mu_T/2 - 2\mu_T$), charm quark mass ($1.45 - 1.55$ GeV), respectively, where $\mu_T = (4m_c^2 + p_T^2)^{1/2}$. We also take into account the uncertainties in the ME's, $\langle O_8(^1S_0)\rangle$ and $\langle O_8(^3P_0)\rangle$ [10].

Let us compare our results for α with the CDF data. We present our result in the form of an error band obtained by combining in quadrature all the theoretical errors described previously. In Fig. 1(a), we compare our result for direct ψ' as a function of p_T with the CDF data[4]. But the error bars in the CDF data are too large to draw any definitive conclusions. We next consider the polarization variable α for prompt J/ψ. The method for calculating cascade effect in prompt J/ψ polarization is explained in Ref. [10]. In Fig. 1(b), we compare our result for α as a function of p_T with the CDF data [4]. Our result for α is small around $p_T = 5$ GeV, but it increases with p_T. Our result is in good agreement with the CDF measurement at intermediate values of p_T, but it disagrees in the highest p_T bin, where the CDF measurement is consistent with 0. The solid lines in Fig. 1(b) are the central curves of α for direct J/ψ and for J/ψ from χ_c. The α for direct J/ψ is smaller than that for direct ψ', because the ME's are different [10]. In the moderate p_T region, the contributions from ψ' and from χ_c add to give an increase in the transverse polarization of prompt J/ψ compared to direct J/ψ. In the high p_T region, the contributions from ψ' and χ_c tend to cancel.

The CDF measurement of the polarization of prompt J/ψ presents a serious challenge to the NRQCD factorization formalism for inclusive quarkonium production. The qualitative prediction that α should increase at large p_T seems inescapable. However, it is still worthwhile to investigate the NLO effect in the fragmentation processes, since the dramatic discrepancy appears at large p_T. Recently, Braaten and Lee performed the full NLO calculation of the color-octet $^3S_1$ gluon fragmentation function for polarized heavy quarkonium [16]. It is not sufficient to get NLO prediction of α, but one can first study the production rate at large p_T in NLO accuracy.

In Run II of the Tevatron, the data sample for J/ψ should be more than an order of magnitude larger than in Run I, which will allow both the production



Figure 1: Polarization variable α vs. p_T for (a) direct ψ' and (b) prompt J/ψ compared to CDF data.

rate and the polarization to be measured with higher precision and out to larger values of p_T. If the result continues to disagree with the predictions of the NRQCD factorization approach, it would indicate a serious flaw in our understanding of inclusive charmonium production. The predictions of low-order perturbative QCD for the spin-dependence of $c\bar{c}$ cross sections could be wrong, or the use of NRQCD to understand the systematics of the formation of charmonium from the $c\bar{c}$ pair could be flawed, or m_c could simply be too small to apply the factorization approach to the charmonium system.

Acknowledgments

The author thanks Eric Braaten and Bernd A. Kniehl for their enjoyable collaboration on the subject discussed here. This work was supported in part by the Alexander von Humboldt Foundation.

References

1. G.T. Bodwin, E. Braaten, and G.P. Lepage, *Phys. Rev.* D **51**, 1125 (1995); **55**, 5855(E) (1997).
2. P. Cho and M.B. Wise, *Phys. Lett.* B **346**, 129 (1995).
3. E. Braaten and S. Fleming, *Phys. Rev. Lett.* **74**, 3327 (1995).
4. CDF Collaboration, hep-ex/0004027.
5. M. Beneke and I.Z. Rothstein, *Phys. Lett.* B **372**, 157 (1996); **389**, 769(E) (1996).
6. M. Beneke and M. Krämer, *Phys. Rev.* D **55**, 5269 (1997).
7. A. Leibovich, *Phys. Rev.* D **56**, 4412 (1997).
8. CDF Collaboration, *Phys. Rev. Lett.* **79**, 578 (1997).
9. CDF Collaboration, CDF Note 3121.
10. E. Braaten, B. Kniehl and J. Lee, hep-ph/9911436.
11. P. Cho and A. Leibovich, *Phys. Rev.* D **53**, 150 (1996); *ibid.* **53**, 6203 (1996).
12. F. Maltoni, private communication on world average of $\Gamma(\chi_{c2} \to \gamma\gamma)$.
13. CDF Collaboration, *Phys. Rev. Lett.* **79**, 572 (1997).
14. B.A. Kniehl and G. Kramer, *Eur. Phys. J.* **C 6**, 493 (1999); *Phys. Rev.* D **60**, 014006 (1996).
15. A.D. Martin, R.G. Roberts, W.J. Stirling, and R.S. Thorne, *Eur. Phys. J.* **C 4**, 463 (1998); CTEQ Collaboration, hep-ph/9903282.
16. E. Braaten and J. Lee, hep-ph/0004228.